

Handbook of Limited Use Drug Products

September 2003



The following contains a list of Limited Use drugs and reimbursement criteria. The drugs are listed in therapeutic categories by brand name. Drugs with similar criteria are grouped together. An index is provided at the end of the book which provides page numbers and a cross reference of generic to brand names. For the complete list of products and more information on the Limited Use process, please refer to the Ontario Drug Benefit Formulary/Comparative Drug Index.

Table of Contents

2	Alzheimer Drugs
3	Analgesics and Anti-inflammatory Drugs
5	Anticoagulants
6	Anticonvulsant Drugs
8	Anti-infective - Antibiotics
14	Anti-infective - Antifungals
17	Anti-infective - Antivirals
22	Antineoplastic and other Cancer related Drugs
28	Benign Prostatic Hypertrophy (BPH) Drugs
29	Cardiovascular Drugs
34	Dermatology Drugs
35	Diabetes Drugs
36	Gastrointestinal Drugs — Proton-Pump Inhibitors
42	Other Gastrointestinal Drugs
44	Hormone Replacement Drugs
46	Ophthalmic Drugs
49	Osteoporosis Drugs
50	Parkinson's Disease Drugs
52	Respiratory Drugs
57	Rheumatoid Arthritis Drugs
58	Transplant Drugs
59	Miscellaneous Drugs
62	Index

Analgesics and Anti-inflammatory drugs

Drug(s)	LU code	Clinical criteria
Celebrex (celecoxib)	316	<p>Osteoarthritis For patients who have failed an adequate trial of acetaminophen (e.g. acetaminophen 1 g QID for several weeks) and have had:</p> <p>History of a documented, clinically significant ulcer or GI bleed; or Failure or intolerance to at least three listed NSAIDS.</p> <p>NOTE: The maximum daily dose of celecoxib which will be reimbursed for the treatment of osteoarthritis is 200 mg.</p>
	317	<p>Rheumatoid arthritis For patients who have had:</p> <p>History of a documented, clinically significant ulcer or GI bleed; or Failure or intolerance to at least three listed NSAIDS.</p> <p>NOTE: The maximum daily dose of celecoxib which will be reimbursed for the treatment of rheumatoid arthritis is 400 mg.</p>

Analgesics and Anti-inflammatory drugs

Drug(s)	LU code	Clinical criteria
Vioxx (rofecoxib)	318	<p>Osteoarthritis For patients who have failed an adequate trial of acetaminophen (e.g. acetaminophen 1 g QID for several weeks) and have had:</p> <p>History of a documented, clinically significant ulcer or GI bleed; or Failure or intolerance to at least three listed NSAIDS.</p> <p>NOTE: The maximum daily dose of rofecoxib which will be reimbursed for the treatment of osteoarthritis is 25 mg.</p>
Codeine Contin (codeine) Duragesic Transdermal (fentanyl) Oxycontin (oxycodone)	201	For the treatment of pain in patients who cannot tolerate, or have failed treatment with a listed long-acting opioid.
Demerol (meperidine)	270	Limited to 2 weeks supply for acute pain.

Anticoagulants

Drug(s)	LU code	Clinical criteria
Fragmin (dalteparin)	186	For acute treatment of deep venous thrombosis (DVT), for a maximum of three weeks;
Fraxiparine, Fraxiparine Forte (nadroparin)	187	For DVT in pregnant or lactating females;
Innohep (tinzaparin)	188	For DVT in patients whom treatment with warfarin is not tolerated, or contraindicated;
Lovenox (enoxaparin)	189	For DVT in patients who have failed treatment with warfarin.
	323	(For Lovenox and Innohep only): For acute treatment of pulmonary embolism, maximum of three weeks.

Anticonvulsant drugs

Drug(s)	LU code	Clinical criteria
Frisium (clobazam)	23	As adjunctive therapy in the treatment of seizure disorders where control by other listed anticonvulsants has been unsatisfactory.
Lamictal (lamotrigine) Neurontin (gabapentin) Sabril (vigabatrin)	136	As adjunctive therapy in the treatment of seizure disorders where control by other listed anticonvulsants has been unsatisfactory.
Tegretol CR (carbamazepine)	67	For patients who have been tried on conventional carbamazepine with unsatisfactory results due to adverse effects or poor control of symptoms.
Topamax (topiramate)	223	As adjunctive therapy in the treatment of seizure disorders where control by other listed anticonvulsants has been unsatisfactory.

Drug(s)	LU code	Clinical criteria
Topamax Sprinkle cap (topiramate)	321	In children age 16 and under, as adjunctive therapy in the treatment of seizure disorders where control by other listed anticonvulsants has been unsatisfactory.

Anti-infective drugs – Antibiotics

Drug(s)	LU code	Clinical criteria
Avelox (moxifloxacin)	337	For the treatment of patients with: CAP with co-morbidity: Community acquired pneumonia with co-morbid illnesses or failure to first-line therapy.
Levaquin (levofloxacin)		
Tequin (gatifloxacin)	338	COPD with risk: Acute bacterial exacerbation of chronic obstructive pulmonary disease (COPD) with risk factors ¹ ; bronchiectasis. ¹ Risk factors include: poor pulmonary lung function (FEV1 below 50% predicted level), age over 65 years, co-morbid medical illness (congestive heart failure, diabetes, chronic renal failure, chronic liver disease), chronic corticosteroid use, malnutrition, prolonged duration of disease, or 4 or more exacerbations/year.
	339	Step-Down: Step-down therapy after parenteral therapy or hospital/emergency department discharge.
	977	Exceptional cases of allergy or intolerance to all other appropriate therapies.

Drug(s)	LU code	Clinical criteria
Cipro (ciprofloxacin)	332	For the treatment of patients with:
		<p>SST/BJ (Gram negative bacteria): Skin/soft tissue and bone/joint infection due to gram negative bacteria; severe diabetic foot infection; severe otitis externa; decubitus ulcers.</p>
	333	<p>GU Tract: Urinary tract infection/prostatitis/epididymitis caused by (suspected or documented) Pseudomonas; sexually transmitted diseases.</p>
334	<p>COPD with risk: Acute bacterial exacerbation of chronic obstructive pulmonary disease (COPD) with risk factors¹; bronchiectasis; pneumonic illness with cystic fibrosis.</p> <p>¹ Risk factors include: poor pulmonary lung function (FEV1 below 50% predicted level), age over 65 years, co-morbid medical illness (congestive heart failure, diabetes, chronic renal failure, chronic liver disease), chronic corticosteroid use, malnutrition, prolonged duration of disease or 4 or more exacerbations per year.</p>	

Anti-infective drugs – Antibiotics

Drug(s)	LU code	Clinical criteria
<p>(continued) Cipro (ciprofloxacin)</p>	336	<p>Step-Down: Step-down therapy after parenteral therapy or hospital/emergency department discharge; febrile neutropenia.</p>
	350	<p>GI: Traveller's diarrhea; enteric fever syndromes; Crohn's disease.</p>
	353	<p>For the prophylaxis or treatment of <i>B. anthracis</i> exposure.</p>
	977	<p>Exceptional cases of allergy or intolerance to all other appropriate therapies.</p>
<p>Floxin (ofloxacin)</p>	340	<p>For the treatment of patients with: SST/BJ (Gram negative bacteria): Skin/soft tissue and bone/joint infection due to gram negative bacteria; severe diabetic foot infection.</p>
	341	<p>GU Tract: Urinary tract infection/prostatitis/epididymitis; sexually transmitted disease.</p>
	338	<p>COPD with risk: Acute bacterial exacerbation of chronic obstructive pulmonary disease (COPD) with risk factors¹; bronchiectasis.</p>

Drug(s)	LU code	Clinical criteria
<p><i>(continued)</i> Floxin (ofloxacin)</p>		<p>¹ Risk factors include: poor pulmonary lung function (FEV1 below 50% predicted level), age over 65 years, co-morbid medical illness (congestive heart failure, diabetes, chronic renal failure, chronic liver disease), chronic corticosteroid use, malnutrition, prolonged duration of disease, or 4 or more exacerbations per year.</p>
	335	<p>GI: Traveller's diarrhea; enteric fever syndromes.</p>
	339	<p>Step-Down: Step-down therapy after parenteral therapy or hospital/emergency department discharge.</p>
977	<p>Exceptional cases of allergy or intolerance to all other appropriate therapies.</p>	

Anti-infective drugs – Antibiotics

Drug(s)	LU code	Clinical criteria
Fucidin Leo tab (sodium fusidate)	342	As part of combination therapy, for the treatment of serious infections confirmed on culture to be caused by a strain of <i>S. aureus</i> or coagulase-negative staphylococci likely susceptible to fusidic acid where standard anti-staphylococcal agents are precluded because of allergy, resistance or treatment failure.
Mycobutin (rifabutin)	103	For the prevention of Mycobacterium Avium Intracellular (MAI) in: Patients with a CD4+ cell count less than 200/mm ³ with an AIDS defining diagnosis; or
	104	Patients with a CD4+ cell count less than 100/mm ³ without an AIDS-defining diagnosis.

Drug(s)	LU code	Clinical criteria
Zyvoxam (linezolid)	362	For the treatment of patients with:
	363	Methicillin-resistant Staphylococcus species (MRSA, MRSE) infections* in patients who are intolerant or have failed vancomycin therapy, or have contraindications to venous access.
	364	Vancomycin resistant Enterococcus species (VRE) infections* in patients switching from IV linezolid. Step-down therapy for the treatment of methicillin-resistant Staphylococcus species or vancomycin resistant Enterococcus species (VRE) infections* after parenteral therapy or hospital/ emergency department discharge. * Infections must be documented and culture proven. Not approved for colonization (e.g. nares, urine etc). Maximum 28 days supply.

Anti-infective drugs – Antifungals

Drug(s)	LU code	Clinical criteria
Diflucan 10mg/mL oral liquid (fluconazole)	274	<p>For the treatment of oral/esophageal candidiasis in immunocompromised patients (e.g. patients with malignancies and transplant patients) who have failed to respond to nystatin or imidazoles and when oral tablets of fluconazole cannot be tolerated.</p> <p>NETWORK NOTE: For oral candidiasis, network will limit supply to 2 weeks. For esophageal candidiasis, network will limit supply to 6 weeks.</p>
	275	<p>For the treatment of patients with disseminated candidiasis when oral tablets of fluconazole cannot be tolerated.</p> <p>NETWORK NOTE: For disseminated candidiasis, network will limit supply to 6 weeks.</p>
	276	<p>For the treatment of patients with cryptococcal meningitis when oral tablets of fluconazole cannot be tolerated.</p> <p>NETWORK NOTE: For cryptococcal meningitis (initial treatment), network will limit supply to 12 weeks.</p>

Drug(s)	LU code	Clinical criteria
<p><i>(continued)</i> Diflucan 10mg/mL oral liquid (fluconazole)</p>	277	<p>For the treatment of patients with vulvovaginal candidiasis when oral tablets of fluconazole cannot be tolerated.</p> <p>NETWORK NOTE: For vulvovaginal candidiasis, network will limit supply to one dose 150 mg (Repeats no more than every 25 days)</p>
<p>Diflucan tab (fluconazole) 50mg, 100mg</p>	<p>202</p> <p>203</p> <p>204</p> <p>205</p>	<p>For the treatment of thrush in immunocompromised patients (i.e. patients with malignancies and transplant recipients) who are unresponsive to nystatin or imidazole preparations.</p> <p>For the treatment of oroesophageal candidiasis in immunocompromised patients (i.e. patients with malignancies and transplant recipients);</p> <p>For patients with disseminated candidiasis;</p> <p>For the treatment of acute cryptococcal meningitis.</p>

Anti-infective drugs – Antifungals

Drug(s)	LU code	Clinical criteria
Diflucan-150 cap (fluconazole)	235	<p>For the treatment of vaginal candidiasis. Dose: 150mg orally once daily for 1 day.</p> <p>NOTE: Repeats within a 25 day period will not be reimbursed.</p>

Anti-infective drugs – Antivirals

Drug(s)	LU code	Clinical criteria
Cytovene injection (ganciclovir sodium)	12	For the treatment of CMV retinitis secondary to AIDS and other immunosuppressive syndromes.
Famvir (famciclovir)	147	<p>Herpes zoster in patients 50 years of age or older, up to 72 hours* after appearance of lesions. Dose: 500mg 3 times/day for 7 days.</p> <p>* The patient must begin treatment within the time frame specified for the product to be reimbursed. There is no benefit from the therapy begun after this time frame.</p> <p>NETWORK NOTE: Network will limit supply to 7 days and 21 tablets.</p>

Anti-infective drugs – Antivirals

Drug(s)	LU code	Clinical criteria
<p>Tamiflu (oseltamivir phosphate)</p>	371	<p>For the prophylaxis (max: 75mg daily) of institutionalized individuals during confirmed* outbreaks of Influenza A or Influenza B.</p> <p>NOTE: Network will limit supply to 6 weeks.</p>
	372	<p>For the treatment (max: 75 mg twice a day) of institutionalized individuals during confirmed* outbreaks due to: Influenza B or, Influenza A (as an alternative to amantadine) or, Influenza A where new cases have developed despite amantadine prophylaxis.</p> <p>NOTE: Network will limit supply to 5 days.</p> <p>* The outbreak must be confirmed by Public Health.</p>
<p>Valcyte (valganciclovir)</p>	374	<p>For the treatment of CMV retinitis in patients with AIDS.</p>

Drug(s)	LU code	Clinical criteria
Valtrex (valacyclovir)	159	<p>Herpes zoster in patients 50 years of age or older, up to 72 hours* after appearance of lesions. Dose: 1 gram 3 times/day for 7 days.</p> <p>* The patient must begin treatment within the time frame specified for the product to be reimbursed. There is no benefit for the therapy begun after this time frame.</p> <p>NETWORK NOTE: Network will limit supply to 7 days and 42 capsules.</p>

Anti-infective drugs – Antivirals

Drug(s)	LU code	Clinical criteria
Zovirax 800mg tab (acyclovir)		Where specified, treatment must begin within the time frames indicated for the product to be reimbursed. There is no benefit from the therapy begun after these time frames.
	95	Herpes zoster in immunocompetent patients 50 years of age or older, up to 72 hours after appearance of lesions: Dose: 800mg 5 times/day for 7 days.
	96	Herpes zoster ophthalmicus regardless of age, up to 72 hours after appearance of lesions: Dose: 800mg 5 times/day for 7 days.
	97	Herpes zoster in immunocompromised patients regardless of age and time elapsed from onset: Dose: 800mg 5 times/day for 7 days.
	314	Varicella zoster in immunocompetent patients greater than or equal to 12 years of age, up to 24 hours after appearance of lesions: Dose: 20mg/kg/dose (max. 800mg) 4 times/day for 5 days. NETWORK NOTE: Network will limit supply up to 7 days and up to 35 tablets.

Drug(s)	LU code	Clinical criteria
Zovirax injection (acyclovir)		In immunocompromised patients who are not responding to oral acyclovir, famciclovir or valacyclovir, and in whom there is concern about the absorption of oral acyclovir, famciclovir or valacyclovir for the treatment of:
	166	Herpes simplex infection, or
	167	Herpes zoster infection.
		In immunocompromised patients who are unable to take oral acyclovir, famciclovir or valacyclovir, for the treatment of:
	168	Herpes simplex infection, or
	169	Herpes zoster infection.

Antineoplastic and other Cancer-related drugs

Drug(s)	LU code	Clinical criteria
Arimidex (anastrozole) Femara (letrozole)	365	For the treatment of metastatic breast cancer in hormone receptor positive post-menopausal women.
Aromasin (exemestane)	180	For the hormonal treatment of metastatic breast cancer in hormone receptor positive post-menopausal women who have disease progression following tamoxifen therapy.
Temodal (temozolomide)	320	For patients with recurrent or progressive glioblastoma multiforme or anaplastic astrocytoma.
Thyrogen (thyrotropin alfa)	368	For use in the monitoring of patients with well-differentiated thyroid cancer.

Antineoplastic and other Cancer-related drugs

Drug(s)	LU code	Clinical criteria
<p>Xeloda (capecitabine)</p>	346	<p>For the first-line treatment of patients with metastatic colorectal cancer in whom combination chemotherapy is not recommended.</p> <p>NOTE: Not to be used in patients who have failed 5-fluorouracil.</p>
	360	<p>For the treatment of metastatic breast cancer in combination with docetaxel in women who experience disease progression on or after an anthracycline.</p>
<p>Bonefos, Ostac (clodronate)</p>	280	<p>For the control and prophylaxis of hypercalcemia of malignancy.</p>
	358	<p>For the treatment of bony metastases in patients with breast cancer.</p>
	359	<p>For the prevention and treatment of osteolytic lesions in patients with multiple myeloma.</p>
<p>Didronel (etidronate)</p>	237	<p>For the treatment of hypercalcemia of malignancy.</p>

Antineoplastic and other Cancer-related drugs

Drug(s)	LU code	Clinical criteria
Intron-A (interferon alfa-2B)	28	For hairy cell leukemia.
	29	For Kaposi's Sarcoma.
Roferon-A (interferon alfa-2A)		
Wellferon (interferon alfa-N1)		
Leustatin (cladribine)	99	For hairy cell leukemia, as a single 7-day treatment course.
Anzemet (dolasetron)	229	For the treatment of emesis in cancer patients receiving highly emetogenic chemotherapy.
	230	For patients receiving intravenous chemotherapy who have not experienced adequate control with other available anti-emetics.
	231	For patients receiving intravenous chemotherapy who experience intolerable side effects with other anti-emetics.
		NOTE: The therapeutic value of Anzemet more than 24 hours after the last dose of chemotherapy is unproven.

Antineoplastic and other Cancer-related drugs

Drug(s)	LU code	Clinical criteria
Kytril (granisetron)	91	For the treatment of emesis in cancer patients receiving highly emetogenic chemotherapy.
	92	For patients receiving intravenous chemotherapy or radiation therapy who have not experienced adequate control with other available anti-emetics.
	93	For patients receiving intravenous chemotherapy or radiation therapy who experience intolerable side effects with other anti-emetics.
	326	<p>For the treatment of emesis in patients receiving radiation therapy which consists of single fraction treatment to the abdominal cavity, hemi-body irradiation and total body irradiation.</p> <p>NOTE: The therapeutic value of Kytril more than 24 hours after the last dose of chemotherapy is unproven.</p>

Antineoplastic and other Cancer-related drugs

Drug(s)	LU code	Clinical criteria
Zofran (ondansetron)	215	For the treatment of emesis in cancer patients receiving highly emetogenic chemotherapy.
	216	For patients receiving intravenous chemotherapy or radiation therapy who have not experienced adequate control with other available anti-emetics.
	217	For patients receiving intravenous chemotherapy or radiation therapy who experience intolerable side effects with other anti-emetics.
	218	<p>For the treatment of emesis in patients receiving radiation therapy which consists of single fraction treatment to the abdominal cavity, hemi-body irradiation and total body irradiation.</p> <p>NOTE: The therapeutic value of Zofran more than 24 hours after the last dose of chemotherapy is unproven.</p>

Antineoplastic and other Cancer-related drugs

Drug(s)	LU code	Clinical criteria
Marinol (delta-9-tetrahydrocannabinol)	40	For the treatment of emesis associated with cancer chemotherapy in patients who are unresponsive to conventional antiemetic therapy.
Imodium (loperamide) Lomotil (diphenoxylate/atropine)	113	For the treatment of diarrhea associated with cancer, including chemotherapy or radiation therapy.
Tantum (benzydamine)	240	For the symptomatic relief of treatment induced mucositis in cancer patients.

Cardiovascular drugs

Drug(s)	LU code	Clinical criteria
Aggrenox (dipyridamole/ ASA)	349	For the secondary prevention of stroke.
Amatine (midodrine)	01	For the treatment of patients disabled by moderate to severe neurogenic orthostatic hypotension (i.e. drop in systolic BP ≤ 20 mm Hg from supine to standing position), in whom conventional nonpharmacologic and pharmacologic (i.e., fludrocortisone) therapies have proven ineffective or are poorly tolerated.
Coreg (carvedilol)	183	For patients with: <ul style="list-style-type: none"> a) NYHA Class II or III Congestive Heart Failure (CHF); <li style="text-align: center;">and b) Currently being treated with an angiotensin converting enzyme (ACE inhibitor, diuretics with or without digoxin, or previously treated, and failed these agents; <li style="text-align: center;">and c) An ejection fraction $\leq 35\%$; <li style="text-align: center;">and d) At least one episode of symptomatic CHF within a 12 month period while receiving optimal management.

Cardiovascular drugs

Drug(s)	LU code	Clinical criteria
Atacand (candesartan)	160	For the treatment of hypertension in patients who cannot tolerate diuretics and beta-blockers or in whom diuretics and beta-blockers are contraindicated or are not adequately effective; and
Avapro (irbesartan)		
Cozaar (losartan)		For the treatment of hypertension in patients who have experienced adverse effects with ACE inhibitors, e.g. cough.
Diovan (valsartan)		
Micardis (telmisartan)		
Teveten (eprosartan)		

Drug(s)	LU code	Clinical criteria
<p>Atacand Plus (candesartan/ HCTZ)</p> <p>Avalide (irbesartan/ HCTZ)</p> <p>Diovan-HCT (valsartan/ HCTZ)</p> <p>Hyzaar, Hyzaar DS (losartan/HCTZ)</p> <p>Micardis Plus (telmisartan/ HCTZ)</p> <p>(HCTZ: hydro- chlorothiazide)</p>	325	<p>For the treatment of hypertension in patients:</p> <p>Who cannot tolerate beta-blockers or in whom beta-blockers and diuretics alone are not effective; and</p> <p>Who have experienced adverse effects with ACE inhibitors, e.g. cough); and</p> <p>Who require the addition of a diuretic to achieve adequate hypertension control.</p>

Drug(s)	LU code	Clinical criteria
Ticlid (ticlopidine)	219 220 221	<p>Ticlopidine is restricted to patients with transient cerebral ischemia.</p> <p>Ticlopidine will be reimbursed for patients:</p> <p>Who are known to be, or become, intolerant of ASA;</p> <p>Where ASA is contraindicated;</p> <p>Who continue to have TIA or stroke symptoms while being treated with ASA.</p>
Trental (pentoxifylline)	76	<p>For the treatment of patients with critical limb ischemia (with arterial ulcers, gangrene and/or rest pain) and documented arterial vascular disease.</p> <p>NOTE: Limited use form must specify if arterial ulcers, gangrene and/or rest pain are present.</p>

Dermatology drugs

Drug(s)	LU code	Clinical criteria
Dovonex (calcipotriol)	191	For the treatment of psoriasis in patients who have failed topical corticosteroids alone, or are intolerant to topical corticosteroids.
Neoral (cyclosporine)	177	For the treatment of psoriasis in patients who have failed, or are intolerant to, other systemic therapies, including methotrexate, acitretin or PUVA.
Nix 5% dermal cream (permethrin)	311	For the treatment of patients who have failed on a less costly listed alternative.
Stieva-A Vitamin A acid (tretinoin)	269	For the treatment of acne vulgaris.

Diabetes drugs

Drug(s)	LU code	Clinical criteria
Humalog (insulin lispro)	226	For insulin requiring diabetics using multiple doses of regular insulin (greater or equal to 3 injections per day) and experiencing recurrent hypoglycemia or poor glycemic control. (Humalog and NovoRapid only): For insulin requiring diabetics using an insulin pump.
Humalog Mix25 (insulin lispro/insulin lispro protamine)	227	
NovoRapid (insulin aspart)		
Prandase (acarbose)	175	In patients who cannot tolerate or have failed treatment with other oral hypoglycemic agents or in whom other oral hypoglycemic agents are contraindicated;
	176	In patients who require combination therapy with more than one oral hypoglycemic agent to control their serum glucose concentrations.

Gastrointestinal drugs – Proton-Pump Inhibitors (PPI)

Drug(s)	LU code	Clinical criteria
<p><i>(continued)</i> Losec (omeprazole)</p> <p>Pantoloc (pantoprazole)</p> <p>Prevacid (lansoprazole)</p>	<p>293</p>	<p>Mild to Moderate GERD For the second-line treatment of mild to moderate GERD in patients who have not responded to at least a 4-week course of an H₂-receptor antagonist. Maximum duration: 6 months</p> <p>For initial treatment, the treatment period with a proton pump inhibitor may extend for no longer than 6 months, after which the need for continued treatment should be reassessed. Reassessment could include a repeat of a course of an H₂-receptor antagonist, or a procedural assessment of the condition.</p> <p>NETWORK NOTE: Network will verify previous H₂-receptor antagonist use in past year.</p>
	<p>294</p>	<p>GERD Recurrence For the treatment of recurrence, once GERD has been procedurally confirmed or the patient has undergone step-down therapy. Maximum duration: unlimited</p>

Gastrointestinal drugs – Proton-Pump Inhibitors (PPI)

Drug(s)	LU code	Clinical criteria
<p><i>(continued)</i> Losec (omeprazole)</p> <p>Pantoloc (pantoprazole)</p>		<p>NETWORK NOTE: Network will supply to 7 days. Network will verify that retreatments are reimbursed only after a four-week period has elapsed since the end of the previous treatment.</p>
<p>Prevacid (lansoprazole)</p>	297	<p>Complicated Peptic Ulcers: For the treatment of peptic ulcers in patients that have procedurally confirmed ulcer complications (i.e. perforation, obstruction, large ulcers or GI bleed). Maximum duration: unlimited.</p>
	298	<p>NSAID induced Ulcers: a) For the treatment of NSAID induced ulcers where the NSAID must be continued. Maximum duration: unlimited while on NSAID therapy.</p>
	299	<p>b) For the prophylaxis of NSAID induced ulcers in patients who have ulcer complications (i.e. perforation or recurrent GI bleed) or are at high-risk for complications (i.e. cardiovascular disease, current or previous peptic ulcer disease, or coagulopathies). Maximum duration: unlimited</p>

Gastrointestinal drugs – Proton-Pump Inhibitors (PPI)

Drug(s)	LU code	Clinical criteria
<p><i>(continued)</i> Losec (omeprazole)</p> <p>Pantoloc (pantoprazole)</p> <p>Prevacid (lansoprazole)</p>	<p>300</p> <p>301</p> <p>302</p> <p>303</p> <p>304</p> <p>305</p>	<p>Other Criteria</p> <p>For the treatment of procedurally confirmed pathological hypersecretory conditions (e.g. Zollinger-Ellison syndrome).</p> <p>For patients with a known allergy to or intolerance of H₂ –receptor antagonist.</p> <p>For patients with gastroduodenal Crohn’s disease.</p> <p>For patients with short-gut syndrome.</p> <p>For patients with scleroderma.</p> <p>For patients with pancreatitis.</p> <p>Maximum duration: unlimited</p>

Gastrointestinal drugs – Proton-Pump Inhibitors (PPI)

Drug(s)	LU code	Clinical criteria
<p>Hp-PAC (lansoprazole/ clarithromycin/ amoxicillin)</p>	306	<p>a) For the treatment of <i>H. pylori</i>-positive peptic ulcers where <i>H. Pylori</i> is documented, by serology, breath test or endoscopy, for a one-week course. Maximum duration: 7 days</p>
	307	<p>b) For the retreatment of <i>H. pylori</i>-positive peptic ulcers where <i>H. Pylori</i> recurrence or persistence is documented, by breath test or endoscopy, for a one-week course. Maximum duration: 7 days (after a four-week period has elapsed since the end of the previous treatment)</p> <p>Retreatment decisions should be based upon positive symptoms and positive endoscopy or positive urea breath test. Retreatment should not be based on a positive serology test, as serology tests may remain positive indefinitely. An alternative antibiotic regimen is recommended when initial therapy fails due to concerns of antimicrobial resistance.</p> <p>NETWORK NOTE: Network will limit supply to 7 days. Network will verify that retreatments are reimbursed only after a four-week period has elapsed since the end of the previous treatment.</p>

Other Gastrointestinal drugs

Drug(s)	LU code	Clinical criteria
Cotazym Cotazym ECS 4 Cotazym ECS 8 Cotazym ECS 20	124	Replacement therapy for pancreatic insufficiency secondary to pancreatic surgery (resection);
Creon 5 Creon 10 Creon 20 Creon 25	125	Replacement therapy for pancreatic insufficiency due to chronic pancreatitis;
Pancrease Pancrease MT4 Viokase Viokase 16	126	Replacement therapy for pancreatic insufficiency due to carcinoma of the pancreas;
(pancrelipase)	225	(Cotazym and Creon products only): Replacement therapy for pancreatic insufficiency due to cystic fibrosis.

Drug(s)	LU code	Clinical criteria
Imodium (loperamide)		For the treatment of diarrhea associated with:
Lomotil (diphenoxylate/ atropine)	110	An ileostomy or a colostomy;
	111	Bowel resection, including short bowel syndrome;
	112	Inflammatory Bowel Diseases, i.e. Crohn's disease and Ulcerative Colitis;
	113	Cancer, including chemotherapy or radiation therapy;
	114	HIV/AIDS;
	115	Acute diarrhea in patients in congregated housing, i.e. Long Term Care Facilities (LTCF), or for patients receiving Home Care;
	224	Fecal incontinence.

Hormone Replacement drugs

Drug(s)	LU code	Clinical criteria
Climara 50 Climara 100 Estraderm Estradot	17	For patients in whom an oral estrogen product is contraindicated.
Oesclim Vivelle (estradiol 17-B)	18	For patients in whom an oral estrogen product has been tried and caused an adverse effect.
Estalis-Sequi 140/50, 250/50 (estradiol 17-B & norethindrone acetate + estradiol 17-B)	138	For patients in whom a combination of a less costly listed oral estrogen product and oral progestin is contraindicated.
Estalis Transdermal Pad 140/50 mcg 250/50 mcg (norethindrone acetate & estradiol 17-B)	139	For patients in whom a combination of a less costly listed oral estrogen product and oral progestin has been tried and caused an adverse effect.
Estracomb (estraderm 50 & estragest)		

Drug(s)	LU code	Clinical criteria
Estrogel (estradiol 17-B)	308	For patients in whom an oral estrogen product is contraindicated.
	288	For patients in whom an oral estrogen product has been tried and caused an adverse effect. NETWORK NOTE: Network will limit supply to one pump per month.

Ophthalmic drugs

Drug(s)	LU code	Clinical criteria
Alphagan (brimonidine tartrate)	171	As first line treatment of elevated intraocular pressure in patients who cannot tolerate an ophthalmic beta-blocking agent or where beta- blocking agents are contraindicated;
Azopt (brinzolamide)		
Trusopt (dorzolamide)	172	As adjunctive therapy of elevated intraocular pressure in patients who cannot tolerate oral carbonic anhydrase therapy, or topical pilocarpine therapy, or topical beta-blocker therapy, or in whom these therapies have been ineffective.
Travatan (travoprost)		
Xalatan (latanoprost)		
Cosopt (dorzolamide/ timolol maleate)	309	For the treatment of patients with elevated intraocular pressure in whom: First-line agents were not tolerated or are contraindicated (i.e. oral carbonic anhydrase therapy or topical pilocarpine therapy).
	310	First-line agents were ineffective (i.e. oral carbonic anhydrase therapy, topical pilocarpine therapy, or topical beta-blocker therapy alone).
Botox (botulinum toxin type A)	10	For the treatment of strabismus and blepharospasm associated with dystonia, including benign essential blepharospasm or VII nerve disorders in patients 12 years of age or older.

Drug(s)	LU code	Clinical criteria
<p>Duolube, Lacri-Lube (petrolatum/ mineral oil)</p> <p>Hypotears, Liquifilm Tears (polyvinyl alcohol)</p> <p>Isopto Tears 0.5%, 1%, Murocel 1% (methylcellulose)</p> <p>Tears Naturale (dextran 70 & hydroxypropyl methylcellulose)</p> <p>Tears Naturale II (dextran 70 & hydroxypropyl methylcellulose & polyquad)</p> <p>Tears Plus (polyvinyl alcohol & polyvinylpyrrol- idone)</p>	49	For patients with objective evidence of keratoconjunctivitis sicca as confirmed by filamentary keratopathy on slit lamp examination or biopsy.

Ophthalmic drugs

Drug(s)	LU code	Clinical criteria
Ocuflox 0.3% (ofloxacin)	170	For the treatment of conjunctivitis caused by susceptible strain(s) of Staphylococcus aureus , Staphylococcus epidermidis , Streptococcus pneumoniae and Hemophilus influenzae which is/are resistant or unresponsive to listed alternative agents.

Osteoporosis drugs

Drug(s)	LU code	Clinical criteria
Actonel (risedronate) 5mg, 35mg	369	For the treatment of osteoporosis in patients who have:
Fosamax (alendronate) 10mg, 70mg		Two out of the following three criteria: BMD at least 3.0 standard deviations below the young adult mean, age of 75 or greater, prior osteoporosis-related fracture; or
	370	Failed* or , experienced intractable side effects, or have a contraindication to, cyclical etidronate (Didrocal) therapy. * Failure is defined as: continued loss of bone mineral density (loss of more than 3%) after two years of therapy, or a new osteoporosis related fracture after one year of therapy.
Evista (raloxifene)	373	For the treatment of osteoporosis in postmenopausal women who have: Failed* or , experienced intractable side effects, or have a contraindication to, alendronate OR risedronate. * Failure is defined as continued loss of bone mineral density (loss of more than 3%) after two years of therapy, or a new osteoporosis related fracture after one year of therapy.

Parkinson's Disease drugs

Drug(s)	LU code	Clinical criteria
Comtan (entacapone)	367	For the treatment of patients with Parkinson's disease with 25% of the waking day in the off state despite maximally tolerated doses of levodopa.

Drug(s)	LU code	Clinical criteria
Sinemet CR (levodopa/ carbidopa)	64	For patients with Parkinson's disease who have been treated with conventional therapy (Prolopa or conventional Sinemet), and experienced adverse effects related to drug level fluctuations, such as ON/OFF or wearing off phenomena.
	65	For patients presently requiring anti-parkinsonian drug administration (levodopa/carbidopa) more than three times daily.

Respiratory drugs

Drug(s)	LU code	Clinical criteria
Alupent Inh soln* (orciprenaline)	256	Patients who have a tracheostomy;
Atrovent Inh soln* (ipratropium)	257	Patients with cystic fibrosis in whom nebulizer therapy is indicated;
Berotec Inh soln* (fenoterol)	258	Patients with severe mental or physical disabilities;
Combivent Inh soln* (ipratropium/ salbutamol)	259	Patients who have previously used nebulizer therapy within the last 12 month period.
Vaponefrin Inh soln* (racemic epinephrine)		
Ventolin Inh soln* (salbutamol)		

Drug(s)	LU code	Clinical criteria
Isuprel Inh soln* (isoproterenol)	260	Children aged 6 years or less;
	261	Patients who have a tracheostomy;
Pulmicort Nebuamp (budesonide)*	262	Patients with cystic fibrosis in whom nebulizer therapy is indicated;
	263	Patients with severe mental or physical disabilities;
	264	Patients who have previously used nebulizer therapy within the last 12 month period.

Respiratory drugs

Drug(s)	LU code	Clinical criteria
Atrovent Inh soln UDV* (ipratropium)	265	Individuals must have a known hypersensitivity to the preservative in the bulk solution, and have a tracheostomy;
Ventolin Nebules* (salbutamol)	266	Individuals must have a known hypersensitivity to the preservative in the bulk solution, and be patients with cystic fibrosis in whom nebulizer therapy is indicated;
	267	Individuals must have a known hypersensitivity to the preservative in the bulk solution, and have severe mental or physical disabilities;
	268	Patients who have previously used nebulizer therapy within the last 12 month period.
<p>* NOTE (For nebulizer solution products): For the vast majority of patients, a metered dose inhaler is the preferred therapy. Nebulizer therapy will be reimbursed for patients who are unable to use a metered dose inhaler, including an inhaler with a spacer attachment, or a turbuhaler.</p>		

Drug(s)	LU code	Clinical criteria
Berotec Inh Pdr (fenoterol)	08	For patients who have not responded to other less expensive inhaled beta-2 adrenergic agonists.
Advair, Advair Diskus (salmeterol xinafoate & fluticasone propionate) Symbicort Turbuhaler (budesonide & formoterol fumarate dihydrate)	330	For the treatment of asthma in patients who are using optimum anti-inflammatory treatment and are still experiencing breakthrough symptoms.
Foradil (formoterol fumarate) Oxeze (formoterol fumarate dihydrate) Serevent, Serevent Diskus, Serevent Diskhaler Disks (salmeterol xinafoate)	132	For the treatment of asthma in patients who are using optimum anti-inflammatory treatment and are still experiencing breakthrough symptoms. NOTE: This drug is not for relief of acute symptoms.

Respiratory drugs

Drug(s)	LU code	Clinical criteria
Atrovent nasal spray (ipratropium bromide)	03	For the treatment of non-allergic vasomotor rhinitis.
Tilade Inh (nedocromil sodium)	72	As second line therapy for asthmatics who are intolerant of sodium cromoglycate.

Rheumatoid Arthritis drugs

Drug(s)	LU code	Clinical criteria
Arava (leflunomide)	331	For the treatment of rheumatoid arthritis in patients who have failed, or are intolerant to, one or more of the listed Disease-Modifying Anti-Rheumatic Drugs (DMARDS).
Celebrex (celecoxib)	317	<p>Rheumatoid arthritis: For patients who have had history of a documented, clinically significant ulcer or GI bleed, or Failure or intolerance to at least three listed NSAIDS.</p> <p>NOTE: The maximum daily dose of celecoxib which will be reimbursed for the treatment of rheumatoid arthritis is 400mg.</p>
Neoral (cyclosporine)	178	For the treatment of rheumatoid arthritis in patients who have failed, or are intolerant to, other systemic therapies, including Disease-Modifying Anti Rheumatic Drugs (DMARDS).

Transplant drugs

Drug(s)	LU code	Clinical criteria
Cellcept (mycophenolate)	190	For the prophylaxis of organ rejection in patients receiving allogeneic renal, cardiac or hepatic transplants.
Prograf (tacrolimus)	173	For solid organ transplant and bone marrow transplant.

Miscellaneous Drugs

Drug(s)	LU code	Clinical criteria
Actonel (risedronate sodium) 30mg tab	319	For the treatment of Paget's disease.
Botox 100u/vial (botulinum toxin type A)	10	For the treatment of strabismus and blepharospasm associated with dystonia, including benign essential blepharospasm or VII nerve disorders in patients 12 years of age or older.
	130	To reduce the subjective symptoms and objective signs of cervical dystonia (spasmodic torticollis) in adults.
Detrol (tolterodine L-tartrate)	290	For patients with urinary frequency, urgency, or urge incontinence who have failed to respond to behavioural techniques and oxybutynin has been shown to be either ineffective or resulted in unacceptable side effects.
Didronel (etidronate)	236	For the treatment of Paget's disease.
	237	For the management of hypercalcemia of malignancy.
Fluotic (sodium fluoride)	20	For the treatment of otosclerosis
	21	For the treatment of otospongiosis.

Miscellaneous Drugs

Drug(s)	LU code	Clinical criteria
Lactaid (lactase enzyme) tab, 6.5mL oral liquid package, 15.5mL oral liquid package	31	For the management of lactose intolerance which has been confirmed by history or by lactose tolerance test.
Marinol (delta-9-tetrahydrocannabinol)	40	For the treatment of emesis associated with cancer chemotherapy in patients who are unresponsive to conventional antiemetic therapy.
	345	For the treatment of AIDS-related anorexia associated with weight loss and prescription is from a prescriber approved for the Facilitated Access mechanism (see Part VI of the Formulary/ CDI binder).

Drug(s)	LU code	Clinical criteria
Nimotop (nimodipine)	42	As adjunctive therapy to improve the neurologic outcome following subarachnoid haemorrhage during the acute management period (within 4 days of haemorrhage).
	43	As prophylaxis of ischemia if surgery is delayed.
Sibelium (flunarizine HCl)	60	For patients with migraine headaches who have not responded to propranolol.
	61	For patients who have tried propranolol and experienced significant adverse effects.
	62	For patients in whom propranolol is contraindicated. CAUTIONS: Contraindicated in patients with clinical depression and in patients with extrapyramidal disorders.
Wellbutrin SR (bupropion)	315	For the treatment of depression.

Index

- acarbose, 35
See Prandase
- Actonel, 49, 59
- acyclovir, 19, 20, 21
See Zovirax
- Advair, 55
- Aggrenox, 29
- alendronate, 49
See Fosamax
- alfuzosin, 28
See Xatral
- Alphagan, 46
- Alupent, 52
- Amatine, 29
- anastrozole, 22
See Arimidex
- Anzemet, 24
- Arava, 57
- Aricept, 2
- Arimidex, 22
- Aromasin, 22
- Atacand, 30
- Atacand Plus, 31
- Atrovent, 52, 54, 56
- Avalide, 31
- Avapro, 30
- Avelox, 8
- Azopt, 46
- benzylamine, 27
See Tantum
- Berotec, 52, 55
- Bonafos, 23
- Botox, 46, 59
- botulinum, 46, 59
- brimonidine, 46
See Alphagan
- brinzolamide, 46
See Azopt
- budesonide, 53, 55
See Pulmicort, Symbicort
- bupropion, 61
See Wellbutrin SR
- calcipotriol, 34
See Dovonex
- candesartan, 30, 31
See Atacand, Atacand Plus
- capecitabine, 23
See Xeloda
- carbamazepine, 6
See Tegretol CR
- carvedilol, 29
See Coreg
- Celebrex, 3, 57
- celecoxib 3, 57
See Celebrex
- Cellcept, 58
- Cipro, 9, 10
- ciprofloxacin, 9, 10
See Cipro
- cladribine, 24
See Leustatin
- Climara, 44
- clobazam, 6
See Frisium
- clodronate, 23
See Bonafos, Ostac
- clopidogrel, 32
See Plavix
- Codeine Contin, 4
- Combivent, 52
- Comtan, 50
- Coreg, 29
- Cosopt, 46
- Cotazym, 42
- Cozaar, 30
- Creon, 42
- cyclosporine, 34, 57
See Neoral
- Cytovene, 17
- dalteparin, 5
See Fragmin
- delta-9-tetrahydrocannabinol, 27, 60
See Marinol
- Demerol, 4
- Detrol, 59
- Didronel, 23, 59
- Diflucan, 14, 15
- Diflucan-150, 16
- Diovan, 30
- Diovan-HCT, 31
- diphenoxylate/atropine, 27, 43
See Lomotil
- dipyridamole/ASA, 29
See Aggrenox
- dolasetron, 24
See Anzemet

- donepezil, 2
See Aricept
- dorzolamide, 46
See Trusopt
- dorzolamide/timolol maleate, 46
See Cosopt
- Dovonex, 34
- Duragesic, 4
- Duolube, 47
- epinephrine, 52
See Vaponefrin
- eprosartan, 30
See Teveten
- enoxaparin, 5
See Lovenox
- entacapone, 50
See Comtan
- Estalis, 44
- Estalis-Sequi, 44
- Estracomb, 44
- Estraderm, 44
- estraderm 50 & estragest, 44
See Estracomb
- estradiol 17-B, 44, 45
See Climara, Estraderm, Estradot, Estrogele, Oesclim, Vivelle
- estradiol 17-B & norethindrone acetate + estradiol 17-B, 44
See Estalis-Sequi
- Estradot, 44
- Estrogele, 45
- etidronate, 23, 59
See Didronel
- Evista, 49
- Exelon, 2
- exemestane, 22
See Aromasin
- famciclovir, 17
See Famvir
- Famvir, 17
- Femara, 22
- fenoterol, 52, 55
See Berotec
- fentanyl, 4
See Duragesic
- Flomax, 28
- Floxin, 10, 11
- fluconazole, 14, 15, 16
See Diflucan
- flunarizine, 61
See Sibelium
- Fluotic, 59
- Foradil, 55
- formoterol, 55
See Foradil, Oxeze, Symbicort
- Fosamax, 49
- Fragmin, 5
- Fraxiparine, 5
- Fraxiparine Forte, 5
- Frisium, 6
- Fucidin, 12
- furosemide, 32
See Lasix Special
- gabapentin, 6
See Neurontin
- galantamine, 2
See Reminyl
- ganciclovir, 17
See Cytovene
- gatifloxacin, 8
See Tequin
- granisetron, 25
See Kytril
- Hp-PAC, 41
- Humalog, 35
- Humalog Mix25, 35
- Hypotears, 47
- Hyzaar, Hyzaar DS, 31
- Imodium, 27, 43
- Innohep, 5
- insulin, 35
See Humalog, Humalog Mix25, NovoRapid
- interferon alfa-2A, 24
See Roferon-A
- interferon alfa-2B, 24
See Intron-A
- interferon alfa-N1, 24
See Wellferon
- Intron-A, 24
- ipratropium, 52, 54, 56
- irbesartan, 30, 31
See Avalide, Avapro
- isoproterenol, 53
See Isuprel

Index

- Isopto Tears, 47
Isuprel, 53
Kytril, 25
Lactaid, 60
lactase enzyme, 60
See Lactaid
Lamictal, 6
lamotrigine, 6
See Lamictal
lansoprazole, 36, 37, 38, 39, 40
See Prevacid
lansoprazole/
clarithromycin/
amoxicillin, 41
See Hp-PAC
Lasix Special, 32
latanoprost, 46
See Xalatan
leflunomide, 57
See Arava
letrozole, 22
See Femara
Leustatin, 24
Levaquin, 8
levodopa/
carbidopa, 51
See Sinemet CR
levofloxacin, 8
See Levaquin
linezolid, 13
See Zyvoxam
Lomotil, 27, 43
loperamide, 27, 43
See Imodium
losartan, 30, 31
See Cozaar, Hyzaar,
Hyzaar DS
Losec, 36, 37, 38, 39, 40
Lovenox, 5
Marinol, 27, 60
meperidine, 4
See Demerol
Micardis, 30
Micardis Plus, 31
midodrine, 29
See Amatine
moxifloxacin, 8
See Avelox
Mycobutin, 12
mycophenolate, 58
See Cellcept
nadroparin, 5
See Fraxiparine
nedocromil, 56
See Tilade
Neoral, 34, 57
Neurontin, 6
nimodipine, 61
See Nimotop
Nimotop, 61
Nix, 34
norethindrone acetate
& estradiol 17-B, 44
See Estalis
NovoRapid, 35
Ocuflox, 48
Oesclim, 44
ofloxacin, 10, 11, 48
See Floxin, Ocuflox
omeprazole, 36, 37, 38,
39, 40
See Losec
ondansetron, 26
See Zofran
orciprenaline, 52
See Alupent
oseltamivir, 18
See Tamiflu
Ostac, 23
Oxeze, 55
oxycodone, 4
See Oxycontin
Oxycontin, 4
Pancrease, 42
pancrelipase, 42
See Cotazym, Creon,
Pancrease, Viokase
Pantoloc, 36, 37,
38, 39, 40
pantoprazole, 36, 37,
38, 39, 40
See Pantoloc
pentoxifylline, 33
See Trental
permethrin, 34
See Nix
Plavix, 32
Prandase, 35
Prevacid, 36, 37,
38, 39, 40
Prograf, 58
Pulmicort, 53

- racemic epinephrine, 52
See Vaponefrin
- raloxifene, 49
See Evista
- Reminyl, 2
- rifabutin, 12
See Mycobutin
- risedronate, 49, 59
See Actonel
- rivastigmine, 2
See Exelon
- rofecoxib, 4
See Vioxx
- Roferon-A, 24
- Sabril, 6
- salbutamol, 52, 54
See Ventolin
- salmeterol, 55
See Serevent, Advair
- Serevent, 55
- Sibelium, 61
- Sinemet CR, 51
- sodium fluoride, 59
See Fluotic
- sodium fusidate, 12
See Fucidin
- Stieva-A, 34
- Symbicort, 55
- tacrolimus, 58
- Tamiflu, 18
- tamsulosin, 28
See Flomax
- Tantum, 27
- Tears Naturale, 47
- Tears Naturale II, 47
- Tears Plus, 47
- Tegretol CR, 6
- telmisartan, 30, 31
See Micardis, Micardis Plus
- Temodal, 22
- temozolomide, 22
See Temodal
- Tequin, 8
- tetrahydocannabinol, 27, 60
See Marinol
- Teveten, 30
- Thyrogen, 22
- thyrotropin alfa.
See Thyrogen 22
- Ticlid, 33
- ticlopidine, 33
See Ticlid
- Tilade, 56
- tinzaparin, 5
See Innohep
- tolterodine, 59
See Detrol
- Topamax, 6
- Topamax
 Sprinkle cap, 7
- topiramate, 6, 7
See Topamax, Topamax Sprinkle cap
- Travatan, 46
- travoprost, 46
See Travatan
- Trental, 33
- tretinoin, 34
See Stieva-A, Vitamin A acid
- Trusopt, 46
- valacyclovir, 19
See Valtrex
- Valcyte, 18
- valganciclovir, 18
See Valcyte
- valsartan, 30, 31
See Diovan, Diovan-HCT
- Valtrex, 19
- Vaponefrin, 52
- Ventolin, 52, 54
- vigabatrin, 6
See Sabril
- Viokase, 42
- Vioxx, 4
- Vitamin A acid, 34
- Vivelle, 44
- Wellbutrin SR, 61
- Wellferon, 24
- Xalatan, 46
- Xatral, 28
- Xeloda, 23
- Zofran, 26
- Zovirax, 19, 20, 21
- Zyxoxam, 13

For more information, contact the Drug Programs Branch:
Toronto 416-327-8109 or toll free: 1-866-811-9893
TTY 1-800-387-5559
www.health.gov.on.ca